

# Neutrino Cross Sections Past, Present, and Future

NuInt04



Sam Zeller  
Columbia University

**FNAL Wine & Cheese**

June 1, 2007



NUINT'02

- this week Fermilab has been host to NuInt07 workshop
  - 5<sup>th</sup> Int'l Workshop on  $\nu$ -Nucleus Interactions in the Few-GeV Region
  - devoted solely to topic of **low energy neutrino cross sections**
  - hit some highlights ... exciting time!

# Outline

- **Motivation**

- $\nu$  oscillations and open questions in  $\nu$  physics
- what are the  $\nu$  cross sections most interested in?
- provide a little bit of the history

- **Current & Future Experiments**

- **Neutrino Cross Sections Measurements**

- experimental results on the various  $\nu$  reactions
- past, present, and future

# Neutrino Oscillations

$$P_{12} = \sin^2 2\theta \sin^2 \frac{1.27 \Delta m^2 L}{E}$$

- current picture:  
3  $\nu$ 's and 2 mass scales


$\nu_3$  —————

$$\Delta m^2_{\text{ATM}} \sim 10^{-3} \text{ eV}^2$$

$\nu_2$  —————

$$\Delta m^2_{\text{SOL}} \sim 10^{-5} \text{ eV}^2$$

$\nu_1$  —————




- this is the world we currently live in

# Open Questions in $\nu$ Physics

- some big questions that we will be actively trying to answer ...
  - what are the masses of neutrinos?
  - what is the mass hierarchy?
  - more precisely measure remaining  $\nu$  oscillation pars:
    - what are the precise values of  $\Delta m^2$ ,  $\theta_{23}$
    - is  $\theta_{13}$  non-zero?
  - is CP violated in the  $\nu$  sector?
- if want to address these questions, first need to understand how  $\nu$ 's interact with matter ... to estimate signal, backgrounds



# Neutrino Cross Sections



- to see maximum osc effects need to be at “low energy”
- specifically, we care about  $\nu$  interactions in the range from 100's-MeV-10's-GeV

**T2K:**  $E_\nu \sim 0.7 \text{ GeV}$


**NOvA:**  $E_\nu \sim 2 \text{ GeV}$

**Gran Sasso:**  $E_\nu \sim 17 \text{ GeV}$

- will motivate why new  $\nu$  data is needed (3 regions)

**Super-K atmospheric**


# Antineutrino Cross Sections



- need  $\bar{\nu}$ 's to tell you about mass hierarchy and  $\text{CP}$ 
  - will measure  $\text{CP}$  by comparing oscillation probabilities of  $\nu_\mu \rightarrow \nu_e$  and  $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$
  - need control of  $\bar{\nu}/\nu$  ratio to better precision than size of expected asymmetry
  - would prefer not to rely on extrapolation of model predictions


- so we need to know all of these cross sections for  $\bar{\nu}$ 's as well

# Accelerator-Based $\nu$ “Roadmap”



# Past $\nu$ Cross Section Measurements

- most of present knowledge comes from bubble chamber measurements




- early exps at ANL, BNL, FNAL, CERN, Serpukhov, etc.

- ~20-100% errors due to:

- low statistics (100's of events)
- uncertainties in  $\nu$  flux  
(still have to face today)

- in addition to large errors, results sometimes conflicting (some care in interpreting)

- but despite vintage, very important in constraining MCs (in addition to high statistics electron scattering data)




# Current Predictions

(notice some of the dates!)

QE scattering  
Llewellyn-Smith  
(1972)

Fermi gas model  
Smith-Moniz  
(1972)


Single  $\pi$   
Production  
Rein-Sehgal  
(1981, 1983)



DIS  
Bodek-Yang  
(2005)

# Status

- this is where we've been for a little over 2 decades




- $\nu$  interactions have been part of past, but also part of future



# Status

- this is where we've been for a little over 2 decades



?  
=



- $\nu$  interactions have been part of past, but also part of future
- in fact, much has happened recently ...

# New $\sigma_\nu$ Predictions (2006-2007)

- J.A. Caballero *et al.*, nucl-th/0705.1429 (2007)
- K.S. Kim, L.E. Wright, nucl-th/0705.0049 (2007)
- E. Hernandez *et al.*, PL **B647**, 452 (2007)
- C. Giusti *et al.*, nucl-th/0607037 (2006)
- P. Lava *et al.*, PRC **73**, 064605 (2006)
- K.S. Kuzmin *et al.*, Acta Phys. Polon. **B37**, 2337 (2006)
- M.C. Martinez *et al.*, PRC **73**, 024607 (2006)
- N. Jachowicz *et al.*, NP. Proc. Suppl. **155**, 260 (2006)
- M. Valverde *et al.*, PL **B642**, 218 (2006), PL **B638**, 325 (2006)
- A. Butkevich, S. Kulagin, nucl-th/0705.1051 (2007)
- M. Martini *et al.*, Phys. Rev. **C75**, 034604 (2007)
- J.E. Amaro *et al.*, PRC **75**, 034613 (2007)
- O. Benhar *et al.*, nucl-ex/0603029 (2006)
- R. Bradford *et al.*, hep-ex/0602017 (2006)
- J. Nieves *et al.*, Phys. Rev. **C73**, 025504 (2006)
- A. Meucci *et al.*, Acta Phys. Polon, **B27**, 2279 (2006)
- G. Co, ActaPhys.Polon.**B37**, 2235 (2006)

QE

- H. Nakamura *et al.*, hep-ph/0705.3884 (2007)
- E.A. Paschos *et al.*, hep-ph/0704.1991 (2007)
- M. Sajjad *et al.*, NP **A782**, 179 (2007), PRD **75**, 093003 (2007)
- O. Lalakulich, E.A. Paschos *et al.*, Nucl. Proc. Suppl. 159, 133 (2006), PRD **74**, 014009 (2006)
- S. Ahmad *et al.*, PRD **74**, 073008 (2006)
- O. Benhar, D. Meloni, PRL **97**, 192301 (2006)
- O. Buss *et al.*, PRC **74**, 044610 (2006), Eur. Phys. J. **A29**, 189 (2006)
- L. Alvarez-Ruso *et al.*, PRC **75**, 055501 (2007)
- E.A. Paschos, A. Kartavtsev, Nucl. Proc. Suppl. **159**, 203 (2006), PRD **74**, 054007 (2006)
- S.K. Singh *et al.*, PRL **96**, 241802 (2006)
- D. Rein, L.M. Sehgal, hep-ph/0606185 (2006)
- B.Z. Kopeliovich, Nucl. Phys. Proc. Suppl. **139**, 219 (2006)

single

$\pi$

production

a lot of  
theoretical  
activity!

DIS

- S. Kulagin, R. Petti, hep-ph/0703033 (2007)
- O. Lalakulich *et al.*, PRC **75**, 015202 (2007)
- O. Benhar, D. Meloni, hep-ph/0610403 (2006)
- K.S. Kuzmin *et al.*, Phys. Atom. Nucl. **69**, 1857 (2006)
- L. Leitner *et al.*, PRC **73**, 065502 (2006), PRC **74**, 065502 (2006), Int.J.Mod.Phys. **A22**, 416 (2007)



# New Neutrino Data

- new data: order of magnitude higher statistics available now
  - allows us to confront model calculations in new ways

- **present:** new low energy (1-to-10's-of-GeV)  $\nu$  data from:

results right now

- NOMAD (1995 - 1998)
- K2K (1999 - 2004)
- MiniBooNE (2002 - present)
- MINOS (2005 - present)

coming soon


- **near future:** program of dedicated experiments at FNAL

- SciBooNE (2007)
- MINERvA (2009)

- field very active
- cast of characters ...  
(deserve some intro)

# K2K


- designed to confirm atm  $\nu$  oscillations (Super-K)
- KEK, 12 GeV protons
  - $\langle E_\nu \rangle = 1.3 \text{ GeV}$



- not only first LBL exp ...
- first on our list to provide updated low E  $\sigma_\nu$  meas
- suite of **near detectors** measure un-oscillated  $\nu$


# K2K

- 1kton water Cerenkov detector
- SciFi (water target)
- SciBar (carbon target)





- miniature replica of Super-K (1/50<sup>th</sup> scale)

- PMTs arranged on inner wall detect Cerenkov light



# K2K

- 1kton water Cerenkov detector
- SciFi (water target)
- SciBar (carbon target)





- scintillating fiber tracker (SciFi) in between H<sub>2</sub>O tanks

- unlike 1kton, every particle is above Cerenkov threshold, so can see everything

# K2K

- 1kton water Cerenkov detector
- SciFi (water target)
- SciBar (carbon target)





- installed later (2003)
- >14k scintillator strips read out by WLS fibers (same as MINOS scintillator)
- will be showing results from all 3



# MiniBooNE




- single detector, designed to check LSND
- FNAL Booster, 8 GeV protons
  - $\langle E_\nu \rangle = 0.8 \text{ GeV}$



- Cerenkov detector ( $\text{CH}_2$ )
  - 1M  $\nu$  and  $\bar{\nu}$  events
  - record size samples in this energy range

# SciBooNE

- dedicated  $\nu$  cross section experiment
- FNAL Booster, 8 GeV protons
  - $\langle E_\nu \rangle = 0.8 \text{ GeV}$
- fine-grained SciBar shipped to FNAL




- couples well-known beam, detector
- has come a long way in a short time (experiment 1<sup>st</sup> proposed in Nov 2005)


# SciBooNE Developments

- March: 1st cosmic rays in SciBar and MRD
- April: detectors move to enclosure

SciBar/EC




MRD





# SciBooNE Developments


- exciting news: 1<sup>st</sup>  $\nu$ 's observed two days ago (May 30<sup>th</sup>)




(R. Napora, NuInt07)

# MINERvA

- dedicated  $\nu$  cross section experiment
- FNAL MI, 120 GeV protons
- fully active detector will sit directly upstream of MINOS




- solid scintillator strip detector surrounded by sampling detectors



- extruded triangular scintillator strips embedded w/ WLS fibers (similar to SciBar, MINOS)


# MINERvA



- dedicated  $\nu$  cross section experiment
- FNAL MI, 120 GeV protons

- LE:  $E_{\nu}^{\text{peak}} = 3 \text{ GeV}$
- ME:  $E_{\nu}^{\text{peak}} = 7 \text{ GeV}$
- HE:  $E_{\nu}^{\text{peak}} = 12 \text{ GeV}$



(fluxes for different target positions)



- observe  $\nu$  ints over broad E range
- impressive physics program:
  - all channels from QE to DIS
- nuclear targets (He, C, Fe, Pb)
  - allow detailed study of nuclear effects in  $\nu$ 's for the first time
- diverse collaboration (HEP, NP, e<sup>-</sup>)

# MINERvA Developments

- March: granted CD1/2/3a approval
- already built full module prototype  
(test mechanical structure, source tested for uniformity)
- next year: multi-plane tracking prototype
  - ~20 of these modules
  - start taking data w/ prototype next year




- coupled w/ test beam program planned
  - next Fall
  - MINERvA TB detector in M-Test

•  $\nu$  data in 2009!

# Modern Challenges

## (1) predicting incoming neutrino flux

- situation has been greatly improving but still true that ...



- how well we know  $\nu$  flux sets limit on how well we can measure absolute  $\sigma_\nu$

$$N_\nu = \sigma_\nu \times \epsilon_{\text{det}} \times \Phi_\nu$$


in the spirit of a historical review ...  
example: 8 GeV p scattering off Be shows resulting  $\nu_\mu$  fluxes at 550 m

demonstrates level of uncertainty in not-so-distant past

# Modern Challenges

## (1) predicting incoming neutrino flux

- situation has been greatly improving but still true that ...



ex: new HARP measurement  
([hep-ex/0702024](#))

- how well we know  $\nu$  flux sets limit  
on how well we can measure absolute  $\sigma_\nu$

$$N_\nu = \sigma_\nu \times \epsilon_{\text{det}} \times \Phi_\nu$$

- meson production data is crucial  
(HARP, MIPP, E910, SPY, NA49)
- modern goal: 5-10%

## (2) in an energy range where nuclear effects are important



# Nuclear Effects

- modern  $\nu$  osc exps measure  $\nu$  interactions on nuclear targets (O, C, Fe, Ar, Pb, ...) - need to understand  $\nu$  ints on nuclei


## (1) $\nu$ interaction w/ bound nucleon

simplest & most common approach  
Is to use Fermi gas model:

- Pauli blocking
- Fermi motion

will come back to this topic later

(F. Sanchez, K2K)



(nuclear effects largest at low  $E_\nu$ , low  $Q^2$ )


# Nuclear Effects

- modern  $\nu$  osc exps measure  $\nu$  interactions on nuclear targets (O, C, Fe, Ar, Pb, ...) - need to understand  $\nu$  ints on nuclei

## (2) propagation of final state hadrons in nucleus (“final state interactions”)

- pions and nucleons can rescatter

(can change momentum, direction, identity as travel through nucleus)



- calculate probabilities of all these processes w/ MC methods



# Nuclear Effects

- (1)  $\nu$  interaction with bound nucleon
- (2) propagation of hadrons in nucleus

*low energy is tough,  
but necessary  
that we understand*

- impact kinematics, rates, and observed final states
- studied extensively in  $e, \mu$  DIS, but only glanced at over limited kinematic range in early, low stat  $\nu$  experiments
  - GGM, SKAT (propane-freon)
  - CHARM, CHARMII (marble, glass)
  - Serpukov (Al)
  - FNAL (Ne)
- this situation will soon change (**MINERvA!**) - will point out examples


# Low Energy $\nu$ Interactions in Nuclei

- let's get to some physics results ... structure of rest of talk:

(1) quasi-elastic (QE)  
- dominates  $E_\nu \lesssim 1$  GeV


(2) NC, CC  $1\pi$  production

(3) CC inclusive, DIS  
- dominates  $E_\nu \gtrsim 5$  GeV



# Quasi-Elastic (QE) Scattering

Why important?




- **important for  $\nu$  oscillation experiments**



- 2-body so can reconstruct  $E_\nu$ , tag flavor of incoming  $\nu$
- also dominant interaction  $\sim 1$  GeV, so forms signal samples ...


- **$\nu_e$  appearance:**  $\nu_e$  QE main signature

- **$\nu_\mu$  disappearance:** need  $\sigma_{QE}(E_\nu)$   
since will be looking for distortions  
in shape of survival probability as  
a function of energy



# Quasi-Elastic (QE) Scattering

Why important?



- **important for  $\nu$  oscillation experiments**



- 2-body so can reconstruct  $E_\nu$ , tag flavor of incoming  $\nu$
- also dominant interaction  $\sim 1$  GeV, so forms signal samples ...

- **interesting in their own right**

- valuable probe of structure of nucleon
- extract info on **axial-vector form factor** of the nucleon  
(in a way that is difficult to obtain in  $e^-$  scattering, will show new measurements!)

# To Set the Scale: Past QE Data

$\langle E_\nu \rangle$	Exp	Mode	Target	Year	# QE events
0.7 GeV	ANL	$\nu$	$D_2$	1973	166
0.5 GeV	ANL	$\nu$	$D_2$	1977	600
2 GeV	CERN	$\nu$	$C_3H_8$	1979	26
2 GeV	GGM	$\nu$	$C_3H_8$	1979	622
0.5 GeV	ANL	$\nu$	$D_2$	1981	1737
1.6 GeV	BNL	$\nu$	$D_2$	1981	1138
5-200 GeV	FNAL	$\nu$	$D_2$	1983	362
6-7 GeV	SKAT	$\nu$	$CF_3Br$	1988	464
9 GeV	SKAT	$\nu$	$CF_3Br$	1990	540
54 GeV	BEBC	$\nu$	$D_2$	1990	552
1.6 GeV	FNAL	$\nu$	$D_2$	1990	2538
5-7 GeV	SKAT	$\nu$	$CF_3Br$	1992	1465
2 GeV	GGM	$\bar{\nu}$	$C_3H_8CF_3Br$	1979	766
1.3 GeV	BNL	$\bar{\nu}$	$H_2$	1980	13
16 GeV	FNAL	$\bar{\nu}$	$NeH_2$	1984	405
6-7 GeV	SKAT	$\bar{\nu}$	$CF_3Br$	1988	52
1.2 GeV	BNL	$\bar{\nu}$	$CH_2$	1988	2919
9 GeV	SKAT	$\bar{\nu}$	$CF_3Br$	1990	159
5-7 GeV	SKAT	$\bar{\nu}$	$CF_3Br$	1992	256

- multiple expts have made QE measurements (but stopped as moved on to other things)

neutrino  
( $\nu_\mu n \rightarrow \mu^- p$ )

~10,000 events total

antineutrino  
( $\bar{\nu}_\mu p \rightarrow \mu^+ n$ )


~4,600 events total

# QE Cross Section Measurements

- this data establishes what we know about the QE  $\sigma$



- **~15-20% uncertainty**  
(considerable spread for any given  $E_{\nu}$ )
- less well-known in threshold region important for osc exps
- low E data mostly on  $D_2$   
(preferred for their simplicity, but not really relevant for meas want to make now)



# QE Cross Section Prediction

- QE cross section can be written in terms of nucleon FFs:

C.H. Llewellyn Smith, Phys. Rep. 3C, 261 (1972)

$$\langle N' | J_\mu | N \rangle = \bar{u}(N') \left[ \gamma_\mu F_V(q^2) + \frac{i\sigma_{\mu\nu} q^\nu \xi F_V^2(q^2)}{2M} + \gamma_5 \gamma_\mu F_A(q^2) \right] u(N)$$

- form factors describe underlying nucleon structure (and come from two sources)

# QE Cross Section Prediction

- QE cross section can be written in terms of nucleon FFs:

C.H. Llewellyn Smith, Phys. Rep. 3C, 261 (1972)

$$\langle N' | J_\mu | N \rangle = \bar{u}(N') \left[ \gamma_\mu F_V(q^2) + \frac{i\sigma_{\mu\nu} q^\nu \xi F_V^2(q^2)}{2M} + \gamma_5 \gamma_\mu F_A(q^2) \right] u(N)$$

- $F_V(Q^2)$ : **vector** form factor well-determined (electron scattering)


$$F_V(Q^2) \sim \frac{1}{(1+Q^2/M_V^2)^2}$$

- historically, vector form factor assumed to have dipole form
- $F_V$  in  $\nu$  related to  $G^{n,p}_{E,M}$  in  $e^-$



# Vector Form Factors

- from electron scattering, we now know vector form factor is not really dipole
- see some of the largest deviations at high  $Q^2$   
(high  $Q^2$  has been an evolving story in electron sector, will get back to)



plus updated fits soon from  
“BBBA2005” ([hep-ex/0602017](https://arxiv.org/abs/hep-ex/0602017))  
Bradford, Bodek, Budd, Arrington  
and a new publication

J.J. Kelly, PRC70, 068202 (2004)

- this provides vector portion of QE  $\sigma$  used in  $\nu$  simulations

# QE Cross Section Prediction

- QE cross section can be written in terms of nucleon FFs:

C.H. Llewellyn Smith, Phys. Rep. 3C, 261 (1972)

$$\langle N' | J_\mu | N \rangle = \bar{u}(N') \left[ \gamma_\mu F_V(q^2) + \frac{i\sigma_{\mu\nu} q^\nu \xi F_V^2(q^2)}{2M} + \gamma_5 \gamma_\mu F_A(q^2) \right] u(N)$$


- $F_A(Q^2)$ : **axial-vector** form factor less well known ( $\nu$  scattering)

$$F_A(Q^2) = \frac{g_A}{(1+Q^2/M_A^2)^2}$$

function of a  
single parameter:  
“**axial mass**”

- + nuclear effects (most  $\nu$  exps use relativistic Fermi Gas model)

# Past Determinations of $M_A$



- this parameter must be determined experimentally

- collection of past  $\nu$  data:

$$M_A = 1.03 \text{ GeV}$$


- so, have been using world average largely set by  $D_2$  in our  $\nu$  simulations

- a lot of interest in this & attempts to remeasure

world avg:  $M_A = 1.03 \pm 0.02 \text{ GeV}$   
J.Phys.G28, R1 (2002)

# How Do You Measure $M_A$ ?

(R. Gran, Nu06)



- varying  $M_A$  has two effects:

- (1) **changes total  $\sigma$**   
( $\uparrow M_A, \uparrow \sigma$ )
- (2) **changes  $Q^2$  dependence**  
( $\uparrow M_A \rightarrow \text{harder } Q^2$ )

- typically  $\nu$  experiments measure  $M_A$  using 2<sup>nd</sup> method (many past experiments did both)

# Modern $M_A$ Analyses



## K2K

+ fine-grained detection  
reconstruct energy & direction  
of outgoing proton

## MiniBooNE

+ statistics  
(order magnitude more QE events than  
has been previously available at low E)

$$\nu_\mu n \rightarrow \mu^- p$$



(single ring topology)




# Two Different Approaches

K2K	MiniBooNE
fine-grained detectors (H <sub>2</sub> O SciFi, <sup>12</sup> C SciBar)	Cerenkov detector ( <sup>12</sup> C)
fit three CC $\nu_\mu$ samples: - 1-track : 8555 events - 2-track QE: 2187 events - 2-track non-QE: 2190 events	fit single $\nu_\mu$ QE sample - 193,709 events
max likelihood fit to $Q^2$ & $E_\nu$ - $M_A$ , norm, $\Phi(E_\nu)$ , non-QE/QE, 2-track/1-track, p re-scattering	least-squares fit to $Q^2$ - $M_A$ , Pauli-blocking par
$Q^2 > 0.2 \text{ GeV}^2$	$0 < Q^2 < 1 \text{ GeV}^2$
reconstruct proton	tag muon-only



# K2K QE Selection

- once 2-track events, kinematic info can be used to enhance fraction of QE evts:




(F. Sanchez, NuInt07)

(R. Gran *et al.*, PRD74, 052002 (2006))


# K2K $M_A$ Fit Results

**SciFi,  $H_2O$  ( $Q^2 > 0.2$ ):**

- $M_A = 1.20 \pm 0.12$  GeV
- $\chi^2/df = 261/235$



1<sup>st</sup> determination  
of  $M_A$  on a  
water target!




# K2K $M_A$ Fit Results

NEW!


**SciFi,  $H_2O$  ( $Q^2 > 0.2$ ):**

- $M_A = 1.20 \pm 0.12$  GeV
- $\chi^2/df = 261/235$




**SciBar,  $^{12}C$  ( $Q^2 > 0.2$ ):**

- $M_A = 1.14 \pm 0.11$  GeV
- $\chi^2/df = 118.7/105$



1<sup>st</sup> determination  
of  $M_A$  on a  
water target!




(F. Sanchez, NuInt07)


10% measurements

# MiniBooNE QE Selection


- measure visible  $T_\mu$ ,  $\theta_\mu$  from Cerenkov light produced by  $\mu$



(T. Katori, NuInt07)



(T. Katori, NuInt07)




- events tagged by  $\mu^- \rightarrow e^-$  decay, not by reconstructing proton
- successful because of Booster  $\nu$  flux (starting with lower non-QE bkg)

# MiniBooNE $M_A$ Fit Results

**MiniBooNE,  $^{12}\text{C}$  ( $Q^2 > 0$ ):**

- $M_A = 1.23 \pm 0.20 \text{ GeV}$
- $\kappa = 1.019 \pm 0.011$
- $\chi^2/\text{df} = 32.8/30$

NEW!



by fitting  $Q^2$  distribution in the data, which is sensitive to  $M_A$



# MiniBooNE $M_A$ Fit Results

**MiniBooNE,  $^{12}\text{C}$  ( $Q^2 > 0$ ):**

- $M_A = 1.23 \pm 0.20 \text{ GeV}$

- $\kappa = 1.019 \pm 0.011$

- $\chi^2/\text{df} = 32.8/30$






# MiniBooNE $M_A$ Fit Results

## MiniBooNE, $^{12}\text{C}$ ( $Q^2 > 0$ ):

- $M_A = 1.23 \pm 0.20 \text{ GeV}$
- $\kappa = 1.019 \pm 0.011$
- $\chi^2/\text{df} = 32.8/30$

NEW!



## MiniBooNE, $^{12}\text{C}$ ( $Q^2 > 0.25$ ):

- $M_A = 1.25 \pm 0.12 \text{ GeV}$
- $\chi^2/\text{df} = 26.6/22$

more standard  
approach  
(10% measurement)

# MiniBooNE $M_A$ Fit Results

## MiniBooNE, $^{12}\text{C}$ ( $Q^2 > 0$ ):

- $M_A = 1.23 \pm 0.20 \text{ GeV}$
- $\kappa = 1.019 \pm 0.011$
- $\chi^2/\text{df} = 32.8/30$

NEW!

## MiniBooNE, $^{12}\text{C}$ ( $Q^2 > 0.25$ ):

- $M_A = 1.25 \pm 0.12 \text{ GeV}$
- $\chi^2/\text{df} = 26.6/22$

publication on these results  
will be available in a few days  
(keep an eye out)



important for  
MiniBooNE  $\nu_\mu \rightarrow \nu_e$   
oscillation analysis

want to be able to  
describe QE events  
throughout kinematic space

# MiniBooNE Verification


- one of advantages of having high stats is can plot in 2D  
(197k  $\nu_\mu$  QE events)

data/MC agreement in  
2D after fitting in 1D




(T. Katori, NuInt07)

# Why is this Interesting?




world avg:  $M_A = 1.03 \pm 0.02$  GeV  
J.Phys.G28, R1 (2002)

- **K2K SciFi** ( $\text{H}_2\text{O}$ ,  $Q^2 > 0.2$ )  
Phys. Rev. **D74**, 052002 (2006)  
 $M_A = 1.20 \pm 0.12$  GeV
- **K2K SciBar** ( $^{12}\text{C}$ ,  $Q^2 > 0.2$ )  
 $M_A = 1.14 \pm 0.11$  GeV
- **MiniBooNE** ( $^{12}\text{C}$ ,  $Q^2 > 0.25$ )  
 $M_A = 1.25 \pm 0.12$  GeV
- new results consistent
- 10% measurements of  $M_A$
- modern data measuring systematically higher  $M_A$  (measuring an “effective  $M_A$ ”)



# MiniBooNE Antineutrino QE Data




NEW!

- MiniBooNE switched to  $\bar{\nu}$  running in Jan 2006
- 8,772  $\bar{\nu}_\mu$  QE events
- expect  $\sim \times 2$  more by Aug
- comforting to see that  $\bar{\nu}$  mode pars seem to successfully describe the MiniBooNE  $\bar{\nu}$  data (if didn't get this ... trouble)

# QE Scattering at NOMAD

- **~10,000 candidate  $\nu_\mu$  QE events from 2-track sample**
  - 70% QE purity, 28% efficiency



$$\nu_\mu n \rightarrow \mu^- p$$


(R. Petti, NOMAD)

- **can measure QE  $\sigma$  on  $^{12}\text{C}$  from  $3 < E_\nu < 100 \text{ GeV}$** 
  - higher energy reach than K2K, MiniBooNE
  - **might expect lower  $M_A$ ?! worth keeping an eye on**
  - publication on final results expected in next few months



# QE Events at MINOS

- expects  $\sim 800,000$   $\nu_\mu$  QE events on iron
  - predict  $\sim 70\%$  QE purity
  - will be interesting to see if can extract  $M_A$  from this data (1<sup>st</sup> on Fe)




exciting!

(R. Gran, D. Bhattacharya, NuInt07)

# QE Scattering at MINERvA


- expect  $\sim 800,000$   $\nu_\mu$  QE events in carbon (4 year run), up to 20 GeV
  - high efficiency: 74%
  - and purity: 77%
- plus, event samples on nuclear targets C, Fe, Pb



example of a simulated  
QE event in MINERvA detector

# MINERvA QE Measurements

- very precise determination of axial form factor  $F_A(Q^2)$






$$F_A(Q^2) = \frac{g_A}{(1 + Q^2/M_A^2)^2}$$

- is the axial form factor really dipole?
- access high  $Q^2$  region
- study medium effects

really important to settle this!

# NC Elastic Scattering




- only existing measurement of absolute  $\sigma$



- BNL E734 measured NC elastic scattering  $\sigma$  as a function of  $Q^2$
- how does this compare against modern data?

# MiniBooNE NC Elastic Scattering

- can select an 84% pure NC EL sample (n+p)



C. Cox, NuInt07

- ~10% of total sample
- 1<sup>st</sup> differential  $\sigma$  results from MiniBooNE
- yesterday was 1<sup>st</sup> time shown outside the collab

# Low Energy Neutrino Interactions


- moving up in energy ...

(1) quasi-elastic (QE)  
- dominates  $E_\nu \lesssim 1$  GeV

(2) NC, CC  $1\pi$  production

- NC  $\pi^0$  (res+coh)
- CC  $\pi^+$  (res+coh)
- CC  $\pi^0$  (res)


(3) CC inclusive, DIS  
- dominates  $E_\nu \gtrsim 5$  GeV







# NC $1\pi^0$ Production

Why important?



$$\nu_\mu N \rightarrow \nu_\mu N \pi^0$$


- **important for neutrino oscillation experiments**
  - poses a serious background to  $\nu_\mu \rightarrow \nu_e$  appearance searches,  $\theta_{13}$  (can look electron-like; in past, primary method for estimating = MC)



- have to control  $\pi^0$ s very well since directly impacts sensitivity

# NC $1\pi^0$ Production

Why important?




$$\nu_\mu N \rightarrow \nu_\mu N \pi^0$$

- **important for neutrino oscillation experiments**
  - poses a serious background to  $\nu_\mu \rightarrow \nu_e$  appearance searches,  $\theta_{13}$  (can look electron-like; in past, primary method for estimating = MC)
- **interesting on their own**
  - $\pi^0$  data can tell us about mode of production (resonant or coherent)

# NC $1\pi^0$ Production

resonant  $\pi^0$  production (dominant)

coherent  $\pi^0$  production




- $\Delta$ ,  $N^*$  baryon resonances, which decay to  $N\pi$
- prominent resonance =  $\Delta(1232)$

# NC $1\pi^0$ Production

resonant  $\pi^0$  production (dominant)

coherent  $\pi^0$  production




- 2 measurements at 2 GeV
- together, < 500 events

(neutrino osc exps  
typically assign uncertainties  
on this  $\sigma$  anywhere in  
25-40% range)

# NC $1\pi^0$ Production

resonant  $\pi^0$  production (dominant)

coherent  $\pi^0$  production



- occurs at ~20% resonant rate
- distinct exp'l signature
- nucleus stays in tact,  
forward emitted  $\pi$ , low  $Q^2$
- 1<sup>st</sup> considered because  
dominant bkg to  $\nu_\mu e \rightarrow \nu_\mu e$


# NC $1\pi^0$ Production

resonant  $\pi^0$  production (dominant)

coherent  $\pi^0$  production



models predict  
wide range

(typically, 100%  
 $\sigma$  uncertainty assumed  
for this process)



(experiments can extract coherent scattering component by looking  
for very forward  $\pi$  production ... will come back to this later in the talk)


# Modern NC $\pi^0$ Analyses

<b>K2K</b>	<b>MiniBooNE</b>
1kton Cerenkov (H <sub>2</sub> O)	800 ton Cerenkov (CH <sub>2</sub> )
$\pi^0$ momentum spectrum	$\pi^0$ momentum spectrum
<b>NC <math>\pi^0</math>/CC <math>\sigma</math> ratio</b>	<b>NC <math>\pi^0</math> COH/RES <math>\sigma</math> ratio</b>
 <p>Diagram of the K2K detector layout showing the Super-Kamiokande detector and the K2K detector. The K2K detector is a cylindrical detector with a central core and an outer shell. The Super-Kamiokande detector is a large cylindrical detector with a central core and an outer shell. The event data shows the following parameters:</p> <pre> NUM 14 RUN 2340 SPIL 41875 SUBRUN 1 EVENT 1368 DATE --Feb- 4 TIME 10:53: 2 TOT PE: 2844.8 MAX PE: 47.9 NPHIT: 467 ANT-PE: 1.3 ANT-MK: 0.6 </pre> <p>TRG ID: 00000001  TMC Ch1: TMC Ch2: TMC  S 21: 0.26: 0.08  T diff: 0.00  Q thr: 0.00  SUB EV: 0/</p>	 <p>Diagram of the MiniBooNE detector layout showing the detector structure and the event data. The event data shows the following parameters:</p> <pre> NUM 14 RUN 2340 SPIL 41875 SUBRUN 1 EVENT 1368 DATE --Feb- 4 TIME 10:53: 2 TOT PE: 2844.8 MAX PE: 47.9 NPHIT: 467 ANT-PE: 1.3 ANT-MK: 0.6 </pre>




# Modern NC $1\pi^0$ Data

- $\pi^0$  mass peak clearly visible
- this is the largest sample of NC  $1\pi^0$  events ever collected!



- 28,600 NC  $\pi^0$  events

J. Link, NuInt07






- 2,496 NC  $\pi^0$  events

S. Nakayama *et al.*, PLB619, 255 (2005)

# $\pi^0$ Momentum Spectra


- modern data revealing some inadequacies in present models
- FSI model?  
(can give effects like this)



# $\pi^0$ Momentum Spectra

- modern data revealing some inadequacies in present models
- FSI model?

Monte Carlo straight out of the box



important for  
MiniBooNE  $\nu_\mu \rightarrow \nu_e$   
oscillation analysis


- extremely important to know how many of these there are
- provided a 10% constraint on NC  $\pi^0$  production rate on  $\text{CH}_2$

# K2K NC $\pi^0$ Cross Section Results

K2K 1kton Water Cerenkov:

$$\text{NC } \pi^0/1\mu \text{ ratio} = 0.064 \\ \pm 0.001 \text{ (stat)} \pm 0.007 \text{ (syst)}$$

- 10% measurement at  $\langle E_\nu \rangle \sim 1.3 \text{ GeV}$
- in good agreement w/ NEUT MC prediction
- 1<sup>st</sup> measurement on a water target



# K2K NC $\pi^0$ Cross Section Results

K2K 1kton Water Cerenkov:

NC  $\pi^0/1\mu$  ratio = 0.064  
 $\pm 0.001$  (stat)  $\pm 0.007$  (syst)

- 10% measurement  
at  $\langle E_\nu \rangle \sim 1.3$  GeV
- in good agreement w/  
NEUT MC prediction
- 1<sup>st</sup> measurement on  
a water target



important for  
Super-K distinguishing

$\nu_\mu \rightarrow \nu_\tau$  VS  $\nu_\mu \rightarrow \nu_s$



# MB NC $\pi^0$ Cross Section Results

## MiniBooNE:

NC  $\pi^0$  COH/RES fraction  
=  $(19.5 \pm 1.1 \text{ (stat)} \pm 2.5 \text{ (syst)})\%$

- 14% measurement at  $\langle E_\nu \rangle \sim 1.1 \text{ GeV}$
- 1<sup>st</sup> measurement of its kind below 2 GeV
- x1.5 lower than default MC prediction (NUANCE, Rein-Sehgal)

(J. Link, NuInt07)




see clear  
evidence  
for NC  
coherent  
 $\pi^0$  prod

# MB NC $\pi^0$ Cross Section Results

## MiniBooNE:

NC  $\pi^0$  COH/RES fraction  
=  $(19.5 \pm 1.1 \text{ (stat)} \pm 2.5 \text{ (syst)})\%$

- 14% measurement  
at  $\langle E_\nu \rangle \sim 1.1 \text{ GeV}$
- 1<sup>st</sup> measurement of its  
kind below 2 GeV
- x1.5 lower than default MC  
prediction (NUANCE, Rein-Sehgal)



important for  
MiniBooNE  $\nu_\mu \rightarrow \nu_e$   
oscillation analysis


recall where we started  
(~100% uncertainties)

not only see in neutrino  
sample, but also in new  
antineutrino data ...




# MiniBooNE Antineutrino $\pi^0$ Data

- with coherent production



- without coherent production




(V. Nguyen, NuInt07)


- ~900 events, enhancement at small angles suggestive of coherent prod

# NC $\pi^0$ Events in MINERvA

- $\pi^0$ 's cleanly identified, 2  $\gamma$ 's tracked
- EM calorimetry is pretty fantastic



measuring  $\pi$  angle important for separating coherent and resonant




- generated & reconstructed angular distributions nearly identical
- highlights MINERvA's excellent  $\pi^0$  angular resolution

# Low Energy Neutrino Interactions



- moving on to the next reaction channel ...

- (1) quasi-elastic (QE)
  - dominates  $E_\nu \lesssim 1$  GeV
- (2) NC, CC  $1\pi$  production
  - NC  $\pi^0$  (res+coh)
  - CC  $\pi^+$  (res+coh)
  - CC  $\pi^0$  (res)
- (3) CC inclusive, DIS
  - dominates  $E_\nu \gtrsim 5$  GeV




# CC $1\pi$ Production

Why important?




- **important for neutrino oscillation analyses**

- poses largest bkg to QE samples ( $\pi$  can be absorbed in nucleus)
- can be a large source of uncertainty for  $\nu_\mu$  disappearance experiments ...
- how well can subtract backgrounds maps into  $\sin^2 2\theta$



# CC $1\pi$ Production

Why important?



$$\nu_\mu N \rightarrow \mu^- N \pi^+$$

- **important for neutrino oscillation analyses**
  - poses largest bkg to QE samples ( $\pi$  can be absorbed in nucleus)
- **interesting on their own**
  - **CC  $\pi^+$** : surpasses QE, largest  $\sigma$  at  $\sim 2$  GeV, so can measure both resonant & coherent  $\pi^+$  with high stats
  - **CC  $\pi^0$** : uniquely probes resonance contribution alone (no coherent contribution in this channel)


# Previous CC $1\pi^+$ $\sigma$ Measurements

- two modes:


$$\nu_{\mu} p \rightarrow \mu^- p \pi^+ (\Delta^{++})$$

$$\nu_{\mu} n \rightarrow \mu^- n \pi^+$$

- no existing data on nuclear targets at low energy ( $E_{\nu} < 3$  GeV)
- total statistics  $\sim 8,000$  events




# K2K CC $1\pi^+$ with SciBar Detector



NEW!

$$\nu_\mu p \rightarrow \mu^- p \pi^+$$

- **1,619 events**
- 41% purity, 13% efficiency



- normalize to QE sample:  
(common approach historically)

$$\nu_\mu n \rightarrow \mu^- p$$

- **8,894 events**
- 60% purity, 60% efficiency

(L. Whitehead, NuInt07)

Sam



# K2K CC $1\pi^+$ /QE $\sigma$ Measurements

K2K SciBar ( $^{12}\text{C}$ ):

NEW!

CC  $1\pi^+p$ /QE  $\sigma = 0.614 \pm 0.061$  (stat)  $+ 0.084$  (nucl)  $+ 0.087$  (syst)  
 $- 0.028$  (nucl)  $- 0.077$  (syst)  
(NEUT prediction = 0.565)



CC  $1\pi^+$ /QE  $\sigma = 0.850 \pm 0.080$  (stat)  $+ 0.127$  (nucl)  $+ 0.119$  (syst)  
 $- 0.039$  (nucl)  $- 0.109$  (syst)  
(NEUT prediction = 0.740)

- ~20% measurements
- in good agreement w/ MC prediction
- consistent w/ MiniBooNE prelim results ([Wascko hep-ex/0602050](http://arxiv.org/abs/hep-ex/0602050))
- measure both total  $\sigma$  and E-dependent  $\sigma$  ...

# K2K CC $1\pi^+$ /QE $\sigma$ Measurement

- in good agreement with past data ...


(L. Whitehead, NuInt07)



- can repeat with same detector in Booster  $\nu$  beam (SciBooNE)
  - higher statistics, lower backgrounds
- measure at MINERvA to extend energy region

# K2K CC $1\pi^+$ /QE $\sigma$ Measurement


- qualitatively good agreement with past data ...



will be important for  
T2K ( $\Delta m^2, \theta_{23}$ )  
goal CC $\pi^+$ /QE: 5-10%

- how well know non-QE backgrounds & their energy dependence will directly map into uncertainties in  $\Delta m^2, \theta_{23}$


# K2K Coherent $1\pi^+$ Production




(M. Hasegawa *et al.*, PRL**95**, 252301 (2005))

- 1<sup>st</sup> CC coh  $\pi^+$  prod results at LE
- somewhat surprising results ...

- see no evidence for coherent  $\pi^+$  prod, **set upper limit**
- MiniBooNE and K2K see evidence for NC coherent  $\pi^0$  prod
- NC, CC difference?  $\mu$  mass effects? (Rein, Sehgal, hep-ph/0606185)




# MINERvA to the Rescue



- coherent  $\pi$  production has been a hot topic (many papers)
- will be able to precisely measure coherent  $\sigma$  as a function of energy
- high statistics
  - 85,000 CC  $\nu + A \rightarrow \mu + A + \pi^+$
  - 37,000 NC  $\nu + A \rightarrow \nu + A + \pi^0$
- compare coherent  $\sigma$  on variety of nuclei (A-dependence, different model predictions for this)

# CC $1\pi^+$ $Q^2$ Dependence?!



(relatively normalized)



(B. Fleming, J. Nowak, NuInt07)

- in addition to coherent  $\pi^+$  prod
- new data revealing large inconsistencies in  $Q^2$  dep of resonance CC  $1\pi^+$  events
  - also seen in other exps
  - and since beginning of NuInt
- **MiniBooNE CC  $1\pi^+$  sample**
  - 71,000 events,  $\text{CH}_2$
  - (order of mag > than all other exp)
- form factors? nuclear effects? coherent  $\pi^+$ ? something else?

# CC $1\pi^+$ Production at MINERvA

- expect 1.2M resonance  $1\pi$  events
- important to be able to track all of the final state particles


(O. Lalakulich)



- can reconstruct hadronic mass (haven't see this yet in modern data sets)
- have been discussing  $\Delta \rightarrow N\pi$  but can also have  $\Delta \rightarrow N\gamma$



# CC $1\pi^+$ Production at MINERvA




important for  
MiniBooNE  $\nu_\mu \rightarrow \nu_e$   
oscillation analysis  
(T2K, NOvA,  $\theta_{13}$ )

- can reconstruct hadronic mass
- opens possibility to measure  $\Delta \rightarrow N\gamma$  in neutrinos for 1<sup>st</sup> time

# Low Energy Neutrino Interactions

- moving on to the next reaction channel ...

- (1) quasi-elastic (QE)
  - dominates  $E_\nu \lesssim 1$  GeV
- (2) NC, CC  $1\pi$  production
  - NC  $\pi^0$  (res+coh)
  - CC  $\pi^+$  (res+coh)
  - CC  $\pi^0$  (res)
- (3) CC inclusive, DIS
  - dominates  $E_\nu \gtrsim 5$  GeV




# Past CC $1\pi^0$ $\sigma$ Measurements

- given  $Q^2$  disagreement and questions surrounding coherent production in CC  $1\pi^+$  channel, can perhaps gain some insight from CC  $\pi^0$  samples (resonance only)


$$\nu_\mu n \rightarrow \mu^- p \pi^0$$

- good example of place where discrepancies exist
- total stats < 2,000 events for all combined




# CC $\pi^0$ Production at K2K

- CC  $\pi^0$  event in the SciBar detector:





(C. Mariani, NuInt07)



- very 1<sup>st</sup> look at this reaction in a modern data set!
- 479 CC  $\pi^0$  events
- 59% purity
- 7.6% efficiency

# K2K CC $\pi^0$ Results

- no coherent scattering component, so probes resonance contrib alone
- venturing into transition region: sample is roughly 1/2 single- $\pi$ , 1/2 multi- $\pi$



# K2K CC $\pi^0$ /QE Cross Section

K2K SciBar:


CC  $\pi^0$ /QE  $\sigma = 0.306 \pm 0.023$  (stat)  $+ 0.023$  (syst)  
 $- 0.021$  (syst)  
(NEUT prediction = 0.220)

NEW!

- ~10% measurement
- 1<sup>st</sup> measurement of CC  $\pi^0$  on carbon at low energy
- 40% higher than MC prediction
- together with agreement in  $1\pi$  samples,  
translates into a  $(30 \pm 18)\%$  excess in multi- $\pi$  cross section



# K2K CC $\pi^0$ /QE Cross Section



- can repeat with same detector in Booster  $\nu$  beam (SciBooNE)
- will be interesting to measure at higher energy (MINERvA)

shouldn't assume we know!



# Low Energy Neutrino Interactions

- and finally ...


(1) quasi-elastic (QE)  
- dominates  $E_\nu \lesssim 1$  GeV

(2) NC, CC  $1\pi$  production

(3) CC inclusive, DIS  
- dominates  $E_\nu \gtrsim 5$  GeV

- “transition region” between QE, DIS  
difficult to predict & poorly measured

- total  $\sigma$  useful as normalization sample for measuring  $\sigma$ 's of other processes






# Inclusive Cross Sections

- MINOS and NOMAD cover interesting region where all three processes contribute (QE, RES, & DIS)

- **MINOS:** 2.1M  $\nu$  Fe  
0.2M  $\bar{\nu}$  Fe (6% WS)

- **NOMAD:** 1.3M events C


- inclusive  $\sigma_\nu$  well-measured at high energy (CCFR, NuTeV)



- less well-known  $E_\nu \approx 3 - 50$  GeV
  - existing data of limited precision ( $\gtrsim 10\%$ )

# Inclusive $\sigma_{CC}$ Measurements


MINOS projection, Fe



(D. Naples, D. Bhattacharya)

- goal: 5% total  $\sigma$  measurement
- $5 < E_\nu < 50$  GeV

NOMAD,  $^{12}\text{C}$ ,  $E_\nu = 20$  GeV





- $6 < E_\nu < 300$  GeV

(R. Petti, NOMAD)

# Inclusive $\sigma_{CC}$ Measurements

MINOS projection, Fe

NOMAD,  $^{12}\text{C}$ ,  $E_\nu = 20 \text{ GeV}$



(R. Petti, NOMAD)

**MINERvA** will bring full power of statistics to bear on this

- millions of events & multiple targets
- 1<sup>st</sup> ever measurement of nuclear effects in  $\nu$ 's!

	Past	Present	Future
QE signal	15-20%	10% $M_A$ - $M_A \sim 1.2$ GeV (nuclear effects?)	
NC $\pi^0$ /QE bkg	25-40%	10% - $\pi^0$ mom differences (FSI?)	
CC $\pi^+$ /QE bkg	25-40%	20% - $Q^2$ disagreement (?)	
CC $\pi^0$ /QE	25-40%	10% - 40% higher than MC (multi- $\pi$ ?)	
coherent $\pi$ bkg	100%	10% NC coh/res $\pi^0$ - tension between NC and CC?	
inclusive	>10%	MINOS, NOMAD will fill in	

	Past	Present	Future
QE signal	15-20%	10% $M_A$ - $M_A \sim 1.2$ GeV (nuclear effects?)	5%
NC $\pi^0$ /QE bkg	25-40%	10% - $\pi^0$ mom differences (FSI?)	<10%
CC $\pi^+$ /QE bkg	25-40%	20% - $Q^2$ disagreement (?)	5%
CC $\pi^0$ /QE	25-40%	10% - 40% higher than MC (multi- $\pi$ ?)	5%
coherent $\pi$ bkg	100%	10% NC coh/res $\pi^0$ - tension between NC and CC?	5-10%
inclusive	>10%	MINOS, NOMAD will fill in	5-10%


+ nuclear effects!

	Past	Present	Future
QE signal	15-20%	10% M - 2 Gr ear effects	5%
NC $\pi^0$ /QE bkg	2%		<10%
CC $\pi^+$ /QE bkg			5%
CC $\pi^0$ /QE			5%
coherent $\pi$ bkg	100%	NC - transition between NC and CC?	5-10%
inclusive	>10%	MINOS, NOMAD will fill in	5-10%

will have to come back to  
this score card in  
a few years to see how  
this all turned out

+ nuclear effects!

# Conclusions

- compelling evidence of  $\nu$  oscillations have increased interest & need for  $\nu$   $\sigma$  measurements
- lot of activity in both exp'ly & theoretically in past few years
  - new results from **K2K, MBooNE, MINOS, NOMAD**
  - 7 new measurements made their debut this week
  - a lot of firsts (many new meas in regions where previously no data)
  - plus, some surprises 
- continued progress on measuring these input  $\sigma$ 's critical for success of future  $\nu$  oscillation experiments
  - **SciBooNE & MINERvA collecting  $\nu$  data soon!**
  - opportunity to make high impact measurements in  $\nu$   $\sigma$ 's

# Special Thanks To

S. Brice  
C. Cox  
R. Gran  
T. Katori  
C. Mariani  
J. Monroe  
J. Morfin  
D. Naples  
J. Nelson  
J. Nowak  
F. Sanchez  
M. Sorel  
R. Petti  
R. Tayloe  
R. VandeWater  
C. Walter  
L. Whitehead